

## 格式

- 向量建议写成  $\boldsymbol{x}_0$

## 内容

- 没有例题吗

# 知识点

## 前言

对于一元函数的极值问题相信大家都十分熟悉，但是对于多元函数的极值问题可能就会比较陌生。大家都学过淑芬怎么可能陌生呢

对于没有限制条件的多元函数来说，只需要对函数求导即可，但是若有了限制条件，即函数的值要在一定条件下才能取到，则需要用到拉格朗日乘子法。

## 引理

设函数  $f(\boldsymbol{x})$

$\varphi_1(\boldsymbol{x}), \varphi_2(\boldsymbol{x}), \dots, \varphi_m(\boldsymbol{x})$  在区域  $D \subset \mathbb{R}^n$  内具有各个连续偏导数，再设  $x_0 = (x_1^0, x_2^0, \dots, x_n^0) \in D$  为  $f(\boldsymbol{x})$  在约束条件  $\begin{cases} \varphi_1(\boldsymbol{x}) = 0 \\ \varphi_2(\boldsymbol{x}) = 0 \\ \vdots \\ \varphi_m(\boldsymbol{x}) = 0 \end{cases}$  下的极值点，并且  $\varphi'(x_0)$  的秩为  $m$  则存在常数  $\lambda_1, \lambda_2, \dots, \lambda_3 \in \mathbb{R}$  使得在  $x_0$  处成立下述等式  $\begin{cases} \frac{\partial f(\boldsymbol{x}_0)}{\partial x_i} + \sum_{j=1}^m \lambda_j \frac{\partial \varphi_j(\boldsymbol{x}_0)}{\partial x_i} = 0 & (i=1, 2, \dots, n) \\ \vdots \\ \frac{\partial f(\boldsymbol{x}_0)}{\partial x_n} + \sum_{j=1}^m \lambda_j \frac{\partial \varphi_j(\boldsymbol{x}_0)}{\partial x_n} = 0 & (j=1, 2, \dots, m) \end{cases}$

## 证明

由于  $\varphi'(\boldsymbol{x}_0)$  的秩为  $m$  我们不妨设行列

式  $\begin{cases} \varphi_1(x_1, x_2, \dots, x_n) \\ \varphi_2(x_1, x_2, \dots, x_n) \\ \vdots \\ \varphi_m(x_1, x_2, \dots, x_n) \end{cases}$  在  $x_0$  处不为零。因此，在  $\boldsymbol{x}_0$  的某个邻域内唯一确定一组具有各个连续偏导数的隐函数  $\begin{cases} x_{n-m+1} = g_1(x_1, x_2, \dots, x_{n-m}) \\ x_{n-m+2} = g_2(x_1, x_2, \dots, x_{n-m}) \\ \vdots \\ x_n = g_m(x_1, x_2, \dots, x_{n-m}) \end{cases}$  满足  $\begin{cases} x_j = g_j(x_1, x_2, \dots, x_{n-m}) & (j=n-m+1, n-m+2, \dots, n) \end{cases}$  且有  $\varphi_k(x_1, \dots, x_{n-m}, g_1(x_1, x_2, \dots, x_{n-m}), g_2(x_1, x_2, \dots, x_{n-m}), \dots, g_m(x_1, x_2, \dots, x_{n-m})) = 0$  将隐函数组代入  $f(\boldsymbol{x}_0)$  得  $f(x_1, \dots, x_{n-m}, g_1(x_1, x_2, \dots, x_{n-m}), g_2(x_1, x_2, \dots, x_{n-m}), \dots, g_m(x_1, x_2, \dots, x_{n-m})) = 0$  因此  $\boldsymbol{x}_0$  是条件极值点转化为  $(x_1^0, x_2^0, \dots, x_{n-m}^0)^0$  为上述函数的通常极值点。

令  $\boldsymbol{x}_0'$  则对  $i=1, 2, \dots, n-m$

有  $\frac{\partial f(\boldsymbol{x}_0)}{\partial x_i} + \frac{\partial f(\boldsymbol{x}_0)}{\partial x_{n-m+1}} + \dots + \frac{\partial f(\boldsymbol{x}_0)}{\partial x_n} = 0$  令  $\frac{\partial f(\boldsymbol{x}_0)}{\partial x_i} = 0$  得  $\frac{\partial f(\boldsymbol{x}_0)}{\partial x_{n-m+1}} + \dots + \frac{\partial f(\boldsymbol{x}_0)}{\partial x_n} = 0$

令

$$\boldsymbol{g}(\boldsymbol{x}) = (g_1(\boldsymbol{x}), g_2(\boldsymbol{x}), \dots, g_m(\boldsymbol{x}))$$

其中  $\boldsymbol{x} = (x_1, x_2, \dots, x_{n-m})$  将上述  $n-m$  个等式写成向量形式，有  $\left( \frac{\partial f(\boldsymbol{x}_0)}{\partial x_1}, \frac{\partial f(\boldsymbol{x}_0)}{\partial x_2}, \dots, \frac{\partial f(\boldsymbol{x}_0)}{\partial x_{n-m}} \right)$   $= \boldsymbol{g}(\boldsymbol{x}_0) = 0$  由于  $\boldsymbol{g}(\boldsymbol{x}_0) = -\left( \begin{array}{c} \frac{\partial f(\boldsymbol{x}_0)}{\partial x_1} \\ \vdots \\ \frac{\partial f(\boldsymbol{x}_0)}{\partial x_{n-m+1}} \end{array} \right)$

由  $\boldsymbol{g}(\boldsymbol{x}_0) = 0$  可得  $\left( \begin{array}{c} \frac{\partial f(\boldsymbol{x}_0)}{\partial x_1} \\ \vdots \\ \frac{\partial f(\boldsymbol{x}_0)}{\partial x_{n-m+1}} \end{array} \right) = -\left( \begin{array}{c} \frac{\partial f(\boldsymbol{x}_0)}{\partial x_1} \\ \vdots \\ \frac{\partial f(\boldsymbol{x}_0)}{\partial x_{n-m+1}} \end{array} \right)$

即  $\left( \begin{array}{c} \frac{\partial f(\boldsymbol{x}_0)}{\partial x_1} \\ \vdots \\ \frac{\partial f(\boldsymbol{x}_0)}{\partial x_{n-m+1}} \end{array} \right) = \left( \begin{array}{c} \frac{\partial f(\boldsymbol{x}_0)}{\partial x_1} \\ \vdots \\ \frac{\partial f(\boldsymbol{x}_0)}{\partial x_{n-m+1}} \end{array} \right)$

注意到  $-A^{-1}B$  是一个  $m \times m$  矩阵，我们可以将其记为  $A^{-1}B$ 。将  $A^{-1}B$  代入之前的式子  $\left( \begin{array}{c} \frac{\partial f(\boldsymbol{x}_0)}{\partial x_1} \\ \vdots \\ \frac{\partial f(\boldsymbol{x}_0)}{\partial x_{n-m+1}} \end{array} \right)$  得

$$\left( \begin{array}{c} \frac{\partial f(\boldsymbol{x}_0)}{\partial x_1} \\ \vdots \\ \frac{\partial f(\boldsymbol{x}_0)}{\partial x_{n-m+1}} \end{array} \right) = \left( \begin{array}{c} \frac{\partial f(\boldsymbol{x}_0)}{\partial x_1} \\ \vdots \\ \frac{\partial f(\boldsymbol{x}_0)}{\partial x_{n-m+1}} \end{array} \right)$$

另外我们可以将  $\left( \begin{array}{c} \frac{\partial f(\boldsymbol{x}_0)}{\partial x_1} \\ \vdots \\ \frac{\partial f(\boldsymbol{x}_0)}{\partial x_{n-m+1}} \end{array} \right)$  改写成  $\left( \begin{array}{c} \frac{\partial f(\boldsymbol{x}_0)}{\partial x_1} \\ \vdots \\ \frac{\partial f(\boldsymbol{x}_0)}{\partial x_{n-m+1}} \end{array} \right) = \left( \begin{array}{c} \frac{\partial f(\boldsymbol{x}_0)}{\partial x_1} \\ \vdots \\ \frac{\partial f(\boldsymbol{x}_0)}{\partial x_{n-m+1}} \end{array} \right)$

\frac{\partial \varphi\_2(\boldsymbol{x}\_0)}{\partial x\_{n-m+2}} \cdots & \cdots \\ \frac{\partial \varphi\_2(\boldsymbol{x}\_0)}{\partial x\_n} \vdots \vdots \ddots \vdots \\ \frac{\partial \varphi\_m(\boldsymbol{x}\_0)}{\partial x\_{n-m+1}} & \cdots \\ \frac{\partial \varphi\_m(\boldsymbol{x}\_0)}{\partial x\_{n-m+2}} \cdots & \cdots \\ \frac{\partial \varphi\_m(\boldsymbol{x}\_0)}{\partial x\_n} \end{array} \right) = 0 \quad \text{tag{5}} \quad \text{将 } \left( \begin{array}{c} \varphi\_4 \\ \varphi\_5 \end{array} \right) \text{ 写成分量形式再加上约束条件即可证明。}

## 拉格朗日乘子法

构造函数  $F(x_1, \dots, x_n, \lambda_1, \dots, \lambda_m) = f(\mathbf{x}) + \sum_{j=1}^m \lambda_j \varphi_j(\mathbf{x})$  则上述求条件极值点的必要条件形式转化为  $F$  的通常极值的必要条件  $\begin{cases} \frac{\partial F(\mathbf{x}_0)}{\partial x_i} = 0 \quad (i=1, 2, \dots, n) \\ \frac{\partial F(\mathbf{x}_0)}{\partial \lambda_j} = 0 \quad (j=1, 2, \dots, m) \end{cases}$  此即拉格朗日乘子法

From:  
<https://wiki.cvbbacm.com/> - CVBB ACM Team



Permanent link:

[https://wiki.cvbbacm.com/doku.php?id=2020-2021:teams:farmer\\_john:2sozx:%E6%95%B0%E5%AD%A6%E7%9F%A5%E8%AF%86%E7%82%B9&rev=1591966081](https://wiki.cvbbacm.com/doku.php?id=2020-2021:teams:farmer_john:2sozx:%E6%95%B0%E5%AD%A6%E7%9F%A5%E8%AF%86%E7%82%B9&rev=1591966081)

Last update: 2020/06/12 20:48