

格式

- 1. 向量建议写成 \mathbf{x}_0

内容

- 1. 没有例题吗

知识点

前言

对于一元函数的极值问题相信大家都十分熟悉，但是对于多元函数的极值问题可能就会比较陌生。大家都学过淑芬怎么可能陌生呢

对于没有限制条件的多元函数来说，只需要对函数求导即可，但是若有了限制条件，即函数的值要在一定条件下才能取到，则需要用到拉格朗日乘子法。

引理

设函数 $f(\mathbf{x})$

$\varphi(\mathbf{x})=(\varphi_1(\mathbf{x}),\varphi_2(\mathbf{x}),\dots,\varphi_m(\mathbf{x}))$ 在区域 $D\subset \mathbb{R}^n (m<n)$ 内具有各个连续偏导数，再设 $\mathbf{x}_0=(x_1^0,x_2^0,\dots,x_n^0)\in D$

为 $f(\mathbf{x})$ 在约束条件 $\begin{cases} \varphi_1(\mathbf{x})=0 \\ \varphi_2(\mathbf{x})=0 \\ \vdots \\ \varphi_m(\mathbf{x})=0 \end{cases}$ 下的极值点，

并且 $\varphi'(\mathbf{x}_0)$ 的秩为 m 则存在常数

$\lambda_1,\lambda_2,\dots,\lambda_m\in\mathbb{R}$ 使得在 \mathbf{x}_0 处成立下述等式 $\begin{cases} \frac{\partial f(\mathbf{x}_0)}{\partial x_i}+\sum_{j=1}^m \lambda_j \frac{\partial \varphi_j(\mathbf{x}_0)}{\partial x_i}=0 \quad (i=1,2,\dots,n) \\ \varphi_j(\mathbf{x}_0)=0 \quad (j=1,2,\dots,m) \end{cases}$

证明

由于 $\varphi'(\mathbf{x}_0)$ 的秩为 m 我们不妨设行列

式 $\frac{\partial(\varphi_1,\varphi_2,\dots,\varphi_m)}{\partial(x_{n-m+1},x_{n-m+2},\dots,x_n)}$ 在 \mathbf{x}_0 处不为零。因此，在 \mathbf{x}_0 的某个邻域内唯一确定一组具有各个连续偏导数的隐函数 $\begin{cases} x_{n-m+1}=g_1(x_1,x_2,\dots,x_{n-m}) \\ x_{n-m+2}=g_2(x_1,x_2,\dots,x_{n-m}) \\ \vdots \\ x_n=g_m(x_1,x_2,\dots,x_{n-m}) \end{cases}$ 满足 $x_j^0=g_j(x_1^0,x_2^0,\dots,x_{n-m}^0) (j=n-m+1,n-m+2,\dots,n)$ 且有 $\varphi_k(x_1,\dots,x_{n-m},g_1(x_1,x_2,\dots,x_{n-m}),\dots,g_m(x_1,x_2,\dots,x_{n-m}))=0$ 将隐函数组代入 $f(\mathbf{x}_0)$

得 $f(x_1,\dots,x_{n-m},g_1(x_1,x_2,\dots,x_{n-m}),\dots,g_m(x_1,x_2,\dots,x_{n-m}))$ 因此 \mathbf{x}_0 是条件极值点转化为 $(x_1^0,x_2^0,\dots,x_{n-m}^0)$ 为上述函数的通常极值点。

令 \mathbf{x}_0 则对 $i=1,2,\dots,n-m$

有 $\frac{\partial f(\mathbf{x}_0)}{\partial x_i}+\frac{\partial f(\mathbf{x}_0)}{\partial x_{n-m+1}}\dots+\frac{\partial f(\mathbf{x}_0)}{\partial x_n}+\frac{\partial f(\mathbf{x}_0)}{\partial x_i}=0$


$\frac{\partial f(\mathbf{x}_0)}{\partial x_i}+\frac{\partial f(\mathbf{x}_0)}{\partial x_{n-m+1}}\dots+\frac{\partial f(\mathbf{x}_0)}{\partial x_n}+\frac{\partial f(\mathbf{x}_0)}{\partial x_i}=0$

令

$$\mathbf{g}(\mathbf{x}) = (g_1(\mathbf{x}), g_2(\mathbf{x}), \dots, g_m(\mathbf{x}))^T$$
 其中 $\mathbf{x} = (x_1, x_2, \dots, x_{n-m})$ 将上述 $n-m$ 个等式写成向量形式，有

$$\left(\frac{\partial f(\mathbf{x}_0)}{\partial x_1}, \dots, \frac{\partial f(\mathbf{x}_0)}{\partial x_{n-m}} \right) + \left(\frac{\partial f(\mathbf{x}_0)}{\partial x_{n-m+1}}, \dots, \frac{\partial f(\mathbf{x}_0)}{\partial x_n} \right) \mathbf{g}(\mathbf{x}_0) = 0$$
 由于 $\mathbf{g}(\mathbf{x}_0) = -\left(\begin{array}{c} \frac{\partial \varphi_1(\mathbf{x}_0)}{\partial x_{n-m+1}} & & \\ \frac{\partial \varphi_1(\mathbf{x}_0)}{\partial x_{n-m+2}} & & \\ \frac{\partial \varphi_1(\mathbf{x}_0)}{\partial x_n} & & \\ \frac{\partial \varphi_2(\mathbf{x}_0)}{\partial x_{n-m+1}} & & \\ \frac{\partial \varphi_2(\mathbf{x}_0)}{\partial x_{n-m+2}} & & \\ \frac{\partial \varphi_2(\mathbf{x}_0)}{\partial x_n} & & \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial \varphi_m(\mathbf{x}_0)}{\partial x_{n-m+1}} & & \\ \frac{\partial \varphi_m(\mathbf{x}_0)}{\partial x_{n-m+2}} & & \\ \frac{\partial \varphi_m(\mathbf{x}_0)}{\partial x_n} & & \end{array} \right)^{-1} \left(\begin{array}{c} \frac{\partial \varphi_1(\mathbf{x}_0)}{\partial x_1} & & \\ \frac{\partial \varphi_1(\mathbf{x}_0)}{\partial x_2} & & \\ \frac{\partial \varphi_1(\mathbf{x}_0)}{\partial x_{n-m}} & & \\ \frac{\partial \varphi_2(\mathbf{x}_0)}{\partial x_1} & & \\ \frac{\partial \varphi_2(\mathbf{x}_0)}{\partial x_2} & & \\ \frac{\partial \varphi_2(\mathbf{x}_0)}{\partial x_{n-m}} & & \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial \varphi_m(\mathbf{x}_0)}{\partial x_1} & & \\ \frac{\partial \varphi_m(\mathbf{x}_0)}{\partial x_2} & & \\ \frac{\partial \varphi_m(\mathbf{x}_0)}{\partial x_{n-m}} & & \end{array} \right) \triangleq -A^{-1}B$
 注意到 $-\left(\frac{\partial f(\mathbf{x}_0)}{\partial x_{n-m+1}}, \dots, \frac{\partial f(\mathbf{x}_0)}{\partial x_n} \right) \cdot A^{-1}$ 是一个 m 维行向量，我们可以将其记为 $-\left(\frac{\partial f(\mathbf{x}_0)}{\partial x_{n-m+1}}, \dots, \frac{\partial f(\mathbf{x}_0)}{\partial x_n} \right) \cdot A^{-1} = \left(\lambda_1, \lambda_2, \dots, \lambda_m \right)$ 将 $\left(\frac{\partial f(\mathbf{x}_0)}{\partial x_1}, \frac{\partial f(\mathbf{x}_0)}{\partial x_2}, \dots, \frac{\partial f(\mathbf{x}_0)}{\partial x_{n-m}} \right) + \left(\lambda_1, \lambda_2, \dots, \lambda_m \right) \left(\begin{array}{c} \frac{\partial \varphi_1(\mathbf{x}_0)}{\partial x_1} & & \\ \frac{\partial \varphi_1(\mathbf{x}_0)}{\partial x_2} & & \\ \frac{\partial \varphi_1(\mathbf{x}_0)}{\partial x_{n-m}} & & \\ \frac{\partial \varphi_2(\mathbf{x}_0)}{\partial x_1} & & \\ \frac{\partial \varphi_2(\mathbf{x}_0)}{\partial x_2} & & \\ \frac{\partial \varphi_2(\mathbf{x}_0)}{\partial x_{n-m}} & & \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial \varphi_m(\mathbf{x}_0)}{\partial x_1} & & \\ \frac{\partial \varphi_m(\mathbf{x}_0)}{\partial x_2} & & \\ \frac{\partial \varphi_m(\mathbf{x}_0)}{\partial x_{n-m}} & & \end{array} \right) = 0$
 另外我们可以将 $\left(\frac{\partial f(\mathbf{x}_0)}{\partial x_{n-m+1}}, \dots, \frac{\partial f(\mathbf{x}_0)}{\partial x_n} \right) + \left(\lambda_1, \lambda_2, \dots, \lambda_m \right) \left(\begin{array}{c} \frac{\partial \varphi_1(\mathbf{x}_0)}{\partial x_{n-m+1}} & & \\ \frac{\partial \varphi_1(\mathbf{x}_0)}{\partial x_{n-m+2}} & & \\ \frac{\partial \varphi_1(\mathbf{x}_0)}{\partial x_n} & & \\ \frac{\partial \varphi_2(\mathbf{x}_0)}{\partial x_{n-m+1}} & & \end{array} \right)$

From:
<https://wiki.cvbbacm.com/> - CVBB ACM Team

Permanent link:
https://wiki.cvbbacm.com/doku.php?id=2020-2021:teams:farmer_john:2sozx:%E6%95%B0%E5%AD%A6:%E7%9F%A5%E8%AF%86%E7%82%B9&rev=1591967909 

Last update: **2020/06/12 21:18**